

Multicurves and regular functions on the representation variety of a surface in $SU(2)$

L. Charles and J. Marché *

Abstract

Given a compact surface Σ , we consider the representation space $\mathcal{M}(\Sigma) = \text{Hom}(\pi_1(\Sigma), SU(2))/SU(2)$. We show that the trace functions associated to multicurves on Σ are linearly independent as functions on $\mathcal{M}(\Sigma)$. The proof relies on the Fourier decomposition of the trace functions with respect to a torus action on $\mathcal{M}(\Sigma)$ associated to a pants decomposition of Σ . Consequently the space of trace functions is isomorphic to the skein algebra at $A = -1$ of the thickened surface.

1 Introduction

Given a compact and oriented surface Σ , one define its representation space as the quotient $\text{Hom}(\pi_1(\Sigma), G)/G$. For $G = \text{Sl}(2, \mathbb{C}), SU(2), \text{Sl}(2, \mathbb{R})$, we obtain three related and celebrated spaces. The first one is an algebraic variety which classifies semi-stable complex bundles of rank 2 over Σ with trivial determinant: it contains the two other ones. The second space is compact and contains an open and dense subset supporting a natural symplectic form. Its quantization provides a construction of a topological quantum field theory (TQFT) which has interesting interactions with the topology of 3-manifolds. Finally the last space contains as a connected component the Teichmüller space, that is, the moduli space of hyperbolic structures on Σ .

The purpose of this article is to study a special class of functions on these spaces called "trace functions". Given a 1-dimensional submanifold γ of Σ and a representation $\rho \in \text{Hom}(\pi_1(\Sigma), G)/G$ one set

$$f_{\gamma, G}([\rho]) = \prod_i (-\text{tr}(\rho(t_i)))$$

*Institut de Mathématiques de Jussieu (UMR 7586), Université Pierre et Marie Curie – Paris 6, Paris, F-75005 France.

where the t_i 's represent the free homotopy classes of the components of γ .

Assume that the Euler characteristic of Σ is negative. We prove that the functions $f_{\gamma,G}$ where γ runs over the isotopy classes of 1-submanifolds of Σ without component bounding a disc are linearly independent as functions on $\text{Hom}(\pi_1(\Sigma), G)/G$ for $G = \text{SU}(2)$ or $\text{Sl}(2, \mathbb{C})$. In the case $G = \text{Sl}(2, \mathbb{C})$, it follows that these functions form a basis of the coordinate ring of the representation variety. Consequently the coordinate ring is isomorphic to the skein algebra of $\Sigma \times [0, 1]$ at $A = -1$, cf. sections 1.1 and 1.2 for precise statements.

D. Bullock showed in [Bul97] that the last assertion is equivalent to proving that the skein algebra has no zero divisors. Bullock's proof rely on a delicate analysis of algebraic relations between trace functions which started in [GM93]. Our strategy is completely different and somewhat simpler: using a pants decomposition of the surface, we define on $\mathcal{M}(\Sigma) = \text{Hom}(\pi_1(\Sigma), \text{SU}(2))/\text{SU}(2)$ an action of the torus $\mathbb{T}^{\mathcal{C}}$ where $\mathcal{C}$ is the set of separating curves of the decomposition. Moreover, 1-dimensional submanifolds of Σ are parametrized up to isotopy by their Dehn coordinates, which is a system of parameters depending on the pants decomposition. Finally, we compute the Fourier decomposition of the trace functions relatively to the action of $\mathbb{T}^{\mathcal{C}}$ and show that one can recover the geometric intersection number of two curves and more generally the full Dehn parameters of a multicurve via its Fourier decomposition. This allows us to prove our assertion.

Our motivation to study the trace functions is the quantization of the representation space $\mathcal{M}(\Sigma)$. First, the space of trace functions is a Poisson algebra, the Poisson bracket being defined with the symplectic structure of Atiyah-Bott [AB83]. Hence the skein algebra at $A = -1$ inherits a Poisson bracket. It appears that the skein algebra at $A = e^{-i\hbar/4}$ is a deformation quantization of this Poisson algebra. This is a consequence of the Goldman formula [Gol86] expressing the bracket of trace function, cf. [BFK99] and [T94].

Not only do we have a formal quantization, but also a strict quantization provided by the topological quantum field theory. Working with the combinatorial version of TQFT [BHMV95] we associate to Σ a family of Hilbert spaces $V_k(\Sigma)$ and to each curve γ a family of operators $(\text{Op}_k(\gamma) : V_k(\Sigma) \rightarrow V_k(\Sigma))$. Then it appears that the natural symbol of this family of operators is the trace function of γ . Indeed, by [MN08] the asymptotic behavior of $\text{Op}_k(\gamma)$ as the level k tends to infinity is controlled at first order by the trace function. Furthermore the composition and the commutator of operators corresponds to the product and Poisson bracket of the trace functions. So the relation between the curve operators and the trace func-

tions is similar to the one in microlocal analysis between operators and their symbol. In this point of view, the interest of our result is to produce non-vanishing trace functions and consequently non-vanishing curve operators. This has non-trivial consequences like the asymptotic faithfulness of the representation of the mapping class group on $V_k(\Sigma)$ provided by TQFT, cf. [And06, FWW02, MN08].

In the next parts of this introduction, we give precise statement of our results (sections 1.1 and 1.2) and on the quantization of $\mathcal{M}(\Sigma)$ (sections 1.3 and 1.4). Note that we do not use skein modules and TQFT in the sequel of the paper. The section 2 introduces the Dehn coordinates for multicurves on a surface while section 3 describe torus actions on $\text{Hom}(\pi_1(\Sigma, \text{SU}(2))/\text{SU}(2)$ which are part of the action-angle coordinates of [JW94]. In Section 4, we give the main ingredients for computing the Fourier coefficients of the trace functions. Section 5 describes the applications of the preceding computations while Section 6 explains the isomorphism of the trace functions algebra with the coordinate ring of the representation variety in $\text{Sl}(2, \mathbb{C})$.

1.1 Results

Let G be the group $\text{SU}(2)$ or $\text{Sl}(2, \mathbb{C})$. Consider the space $\text{Hom}(\pi, G)/G$ of morphisms from a group π to G up to conjugation. For any $t \in \pi$, introduce the function χ_t of $\text{Hom}(\pi, G)/G$ defined by

$$\chi_t([\rho]) = -\text{tr}(\rho(t)), \quad \rho \in \text{Hom}(\pi, G)$$

Let us call a *trace function* any linear combination of the χ_t 's with complex coefficients. Because of the relation

$$\text{tr}(a) \text{tr}(b) = \text{tr}(ab) + \text{tr}(a^{-1}b), \quad \forall a, b \in G \quad (1)$$

the space $\mathcal{T}(\pi, G)$ of trace functions is a subalgebra of the algebra of complex valued functions of $\text{Hom}(\pi, G)/G$.

When π is of finite type, $\mathcal{T}(\pi, \text{Sl}(2, \mathbb{C}))$ is finitely generated and is the coordinate ring of the $\text{Sl}(2, \mathbb{C})$ character variety of π . The relation with the usual definition is explained in part 6.

Assume now that π is the fundamental group of a manifold M . Let us define a trace function $f_{\gamma, G}$ for any isotopy class γ of 1-dimensional compact submanifold of M without arc components. Let $t_1, \dots, t_n$ be the homotopy classes of loops which are freely homotopic to the connected component of a representant of γ and set

$$f_{\gamma, G} = \chi_{t_1} \cdots \chi_{t_n}$$

Since the t_i 's are well-defined up to conjugation and inversion, this product only depends on the isotopy class γ .

We will use these definitions in two cases. First, M is a surface and γ an isotopy class of *multicurve*, that is no component of γ bounds a disc. Second, M is 3-dimensional, the 1-dimensional compact submanifolds without arc component are then called the *links* of M .

Theorem 1.1. *Let G be $SU(2)$ or $Sl(2, \mathbb{C})$ and π be the fundamental group of a compact orientable surface Σ with boundary (possibly empty) and negative Euler characteristic. Then the trace functions $f_{\gamma, G}$, where γ runs over the isotopy classes of multicurves of Σ which do not meet the boundary, form a basis of $\mathcal{T}(\pi, G)$.*

The fact that the trace functions ($f_{\gamma, G}$) are a spanning set follows easily from the relations (1). The proof that they are linearly independent is much more delicate and is the content of theorem 5.3. Let j be the obvious map

$$\text{Hom}(\pi, SU(2))/SU(2) \rightarrow \text{Hom}(\pi, Sl(2, \mathbb{C}))/Sl(2, \mathbb{C}).$$

By the previous theorem, j^* maps $\mathcal{T}(\pi, Sl(2, \mathbb{C}))$ bijectively onto $\mathcal{T}(\pi, SU(2))$.

1.2 The Kauffman module at $A = -1$

Let M be a 3-dimensional oriented compact manifold. The skein module of M at $A = -1$ is defined as the complex vector space freely generated by the set of isotopy classes of links of M quotiented by the relations of Figure 1 for $A = -1$.

We denote by $\mathcal{K}(M, -1)$ this vector space. As a consequence of the relation (1), we have a well-defined map

$$\mathcal{K}(M, -1) \rightarrow \mathcal{T}(\pi, G)$$

sending a link γ to $f_{\gamma, G}$. Here π is the fundamental group of M .

Assume that $M = \Sigma \times [0, 1]$ where Σ is an orientable compact surface. Then by identifying Σ with $\Sigma \times \{1/2\} \subset M$, each multicurve of Σ defines a link in M . By a consequence of the Reidemeister theorem, the family of isotopy classes of multicurves of Σ is a basis of $\mathcal{K}(M, -1)$. As a corollary of Theorem 1.1, we obtain the

Theorem 1.2. *For any compact orientable surface Σ with $\chi(\Sigma) < 0$, the natural map*

$$\mathcal{K}(\Sigma \times [0, 1], -1) \rightarrow \mathcal{T}(\pi, G)$$

is an isomorphism.

Bullock proved in [Bul97] that the kernel of this map consists in the nilradical of $\mathcal{K}(\Sigma \times [0, 1], -1)$. Moreover, it is claimed without proof in [PS00] that this nilradical is zero.

The skein module $\mathcal{K}(\Sigma \times [0, 1], -1)$ is an algebra where the product $\gamma \cdot \delta$ is defined by stacking γ over δ . As a consequence of the relations (1) this product is sent to the product of functions by the isomorphism of Theorem 1.2.

1.3 $\mathcal{T}(\pi, G)$ as a Poisson algebra

In this subsection and the next one, G is the group $\mathrm{SU}(2)$, Σ is a closed orientable surface and π its fundamental group. The space $\mathcal{M} = \mathrm{Hom}(\pi, G)/G$ has a natural topology such that the trace functions are continuous. The subset $\mathcal{M}^s$ consisting of classes of irreducible representations is open and dense. By [AB83] and [Gol84], $\mathcal{M}^s$ is a symplectic manifold. Furthermore Goldman in [Gol86] expressed the Poisson bracket of the trace functions of two curves intersecting transversally as a trace function. Consequently, $\mathcal{T}(\pi, G)$ is a Poisson subalgebra of $\mathcal{C}^\infty(\mathcal{M}^s, \mathbb{C})$. The Poisson bracket appears also naturally on the topological side in the following way.

Let us introduce the Kauffman module $\mathcal{K}(M, -e^{i\hbar/4})$ of a 3-dimensional compact oriented manifold. It is defined as the free $\mathbb{C}[[\hbar]]$ -module generated by the set of isotopy classes of banded links of M quotiented by the relations of Figure 1 with $A = -e^{i\hbar/4}$.




Figure 1: Kauffman relations

One has a natural $\mathbb{C}$ -linear map from $\mathcal{K}(M, -e^{i\hbar/4})$ to $\mathcal{K}(M, -1)$ sending a linear combination $\sum c_i(\hbar)\gamma_i$ of banded links to $\sum c_i(0)\tilde{\gamma}_i$, where $\tilde{\gamma}_i$ is the core of γ_i . The kernel of this map is $\hbar\mathcal{K}(M, -e^{i\hbar/4})$.

Assume now that $M = \Sigma \times [0, 1]$. Using the isomorphism of theorem 1.2, we obtain an exact sequence

$$0 \rightarrow \hbar \mathcal{K}(M, -e^{i\hbar/4}) \rightarrow \mathcal{K}(M, -e^{i\hbar/4}) \xrightarrow{\sigma} \mathcal{T}(\pi, G) \rightarrow 0 \quad (2)$$

Furthermore the multiplication by $\hbar$ is a bijection

$$\mathcal{K}(M, -e^{i\hbar/4}) \rightarrow \hbar \mathcal{K}(M, -e^{i\hbar/4}) \quad (3)$$

The injectiveness follows from the fact that $\mathcal{K}(M, -e^{i\hbar/4})$ is free as a $\mathbb{C}[[\hbar]]$ -module, a basis being given by the family of isotopy classes of multicurves of Σ .

Finally, $\mathcal{K}(M, -e^{i\hbar/4})$ has a natural stacking product $*$ and a natural involution $-$ defined by conjugating coefficients and sending a banded link γ to $S(\gamma)$ where $S : \Sigma \times [0, 1] \rightarrow \Sigma \times [0, 1]$ is defined by $S(x, t) = (x, 1 - t)$. One has $\overline{f * g} = \bar{g} * \bar{f}$.

Theorem 1.3. *Let G be the group $\mathrm{SU}(2)$, Σ a closed surface, π its fundamental group and $M = \Sigma \times [0, 1]$. Then the Kauffman module $\mathcal{K}(M, -e^{i\hbar/4})$ with its product $*$ and involution $-$ is a deformation of the Poisson algebra $\mathcal{T}(\pi, G)$ in the sense that for any $f, g \in \mathcal{K}(M, -e^{i\hbar/4})$ one has*

$$\begin{aligned} \sigma(f * g) &= \sigma(f)\sigma(g), & \sigma(\bar{f}) &= \overline{\sigma(f)} \\ \sigma(\hbar^{-1}(f * g - g * f)) &= \frac{1}{i} \{\sigma(f), \sigma(g)\} \end{aligned}$$

where σ is defined in (2) and $\hbar^{-1}$ is the inverse of the map (3).

The first equation follows from the fact that the isomorphism of theorem 1.2 is an algebra morphism. For the second equation, observe that the classes in $\mathcal{K}(M, -e^{i\hbar/4})$ of the multicurves of Σ are real and that their trace functions are real too. Last formula on the commutator follows from the Goldman formula expressing the Poisson bracket of two trace functions (see [Gol86]). The Poisson bracket depends on the symplectic form which itself depends on a choice of an invariant scalar product on the Lie algebra. We set $\langle A, B \rangle = \mathrm{tr}(AB^*)$ for $A, B \in \mathfrak{su}(2)$. The symplectic form on $\mathrm{Hom}(\pi, G)/G$ we are dealing with is the symplectic reduction of the form $\omega(a, b) = \int_{\Sigma} \langle a \wedge b \rangle$ for $a, b \in \Omega^1(\Sigma, \mathfrak{su}(2))$.

1.4 Topological Quantum Field Theory

Let $\mathbb{C}[A^{\pm 1}]$ be the ring of Laurent polynomials and $\mathcal{K}(M, A)$ be the $\mathbb{C}[A^{\pm 1}]$ -module freely generated by the isotopy classes of banded links in M quotiented by the relations of Figure 1.

By sending $\sum P_i(A)\gamma_i$ to $\sum P_i(-e^{i\hbar/4})\gamma_i$, one identifies the skein module $\mathcal{K}(M, A)$ with a complex subalgebra of $\mathcal{K}(M, -e^{i\hbar/4})$ preserved by the involution $-$.

For any integer $k \geq 2$, the topological quantum field theories (TQFT) coming from Chern-Simons theory with gauge group $SU(2)$ provides a complex Hermitian space $V_k(\Sigma)$ of finite dimension. In the geometric framework, the integer $k - 2$ is called the level while in the combinatorial framework constructed in [BHMV95], the authors use the parameter $p = 2k$. Following them, one has a natural algebra morphism

$$\mathcal{K}(M, \zeta_k) \rightarrow \text{End}(V_k(\Sigma))$$

where $\zeta_k = -e^{i\pi/2k}$ and $\mathcal{K}(M, \zeta_k) := \mathcal{K}(M, A) \otimes_{A=\zeta_k} \mathbb{C}$.

For any $f \in \mathcal{K}(M, A)$ and any $k \geq 2$, we denote by $\text{Op}_k(f)$ the corresponding operator of $V_k(\Sigma)$.

Theorem 1.4. *For any $f, g \in \mathcal{K}(M, A)$, one has*

$$\text{Op}_k(f * g) = \text{Op}_k(f) \text{Op}_k(g), \quad \text{Op}_k(\bar{f}) = \text{Op}_k(f)^*$$

and as $k \rightarrow \infty$

$$\text{tr}(\text{Op}_k(f)) = \left(\frac{k}{4\pi^2}\right)^n \int_{\mathcal{M}^s} \sigma(f) \mu + O(k^{n-1})$$

where n is half the dimension of $\mathcal{M}^s$ and μ is the Liouville measure.

First part follows from the general properties of TQFT. The estimation of the trace has been proved in [MN08]. A factor $4\pi^2$ appears due to the different normalization of the symplectic form. This estimation gives some information on the asymptotic behavior of $\text{Op}_k(f)$.

Define the normalized Hilbert-Schmidt norm of $T \in \text{End}(V_k(\Sigma))$ by

$$\|T\|_{\text{HS}} := (\dim(V_k(\Sigma)))^{-1} \text{tr}(TT^*).$$

Then by theorem 1.3 and 1.4, for any $f \in \mathcal{K}(M, A)$, the Hilbert-Schmidt norm of $\text{Op}_k(f)$ is estimated by the L^2 norm of $g = \sigma(f)$:

$$\|\text{Op}_k(f)\|_{\text{HS}}^2 = (\text{Vol}(\mathcal{M}))^{-1} \int_{\mathcal{M}^s} |g|^2 \mu + O(k^{-1}).$$

The space of sequences $(T_k) \in \prod_{k \geq 2} \text{End}(V_k(\Sigma))$ with a bounded Hilbert-Schmidt norm has a natural filtration $O(0) \supset O(1) \supset O(2) \supset \dots$ where (T_k) is in $O(\ell)$ if $\|T_k\|_{\text{HS}} \leq Ck^{-\ell}$ for some C . It follows from theorems 1.3 and 1.4 that this filtration corresponds to the formal one:

$$(\text{Op}_k(f))_k \in O(\ell) \Leftrightarrow f \in \hbar^\ell \mathcal{K}(M, -e^{i\hbar/4}),$$

for any $f \in \mathcal{K}(M, A)$. In particular, if $f \neq 0$, $\text{Op}_k(f)$ does not vanish when k is sufficiently large.

Using this last observation, one may deduce that the action on $V_k(\Sigma)$ of an homeomorphism of Σ provided by TQFT is asymptotically non trivial as soon as the action of this homeomorphism on the Kauffman module is non trivial, cf. [MN08] for details.

2 Dehn coordinates

Given Σ an oriented compact surface with boundary, we will call *multicurve* a submanifold of Σ which do not meet the boundary and has no component bounding a disc. The Dehn theorem classifies isotopy classes of multicurves by decomposing them into simple pieces: in these simple pieces, we will allow multicurves to meet the boundary transversally as we describe it now.

Let w be the arc of points in S^1 whose angles are in $[-\pi/8, \pi/8]$. For every $m \in \mathbb{N}$ we fix a subset W_m of size m in w invariant by complex conjugation. Let $A = S^1 \times [0, 1]$ be an annulus and introduce a family $A(m, t)$ of multicurves of A indexed by $\mathbb{N} \times \mathbb{Z}$. If m is positive, $A(m, t)$ is the multicurve in A (unique up to isotopy) whose projection on $[0, 1]$ is a submersion, which intersects each boundary circle in W_m and whose algebraic intersection with the curve $\{-1\} \times [0, 1]$ equals t (oriented from 0 to 1). An example is shown in the left hand side of Figure 2. In the case where $m = 0$, $A(m, t)$ consists of t parallel copies of the boundary.

Let T be the surface $\{z \in \mathbb{C}, \text{ s.t. } |z| \leq 1 \text{ and } |z \pm \frac{1}{2}| > \frac{1}{4}\}$. Choose identifications of the boundary circles of T with S^1 such that $1 \in S^1$ is identified respectively with $p_1 = \frac{1}{4}$, $p_2 = \frac{-1}{4}$ and $p_3 = i$. We call T the standard pair of pants (or trinion). Let m_1, m_2, m_3 be three non-negative integers with even sum. Consider a multicurve as in the right hand side of Figure 2 which intersects the boundary circles in $W_{m_1}, W_{m_2}, W_{m_3}$ via the identifications with S^1 . Denote it by $C(m_1, m_2, m_3)$.

Consider now a general surface Σ with negative Euler characteristic. From the classification of surfaces, it appears that Σ is obtained by gluing trinions on boundary components. We will call *pant decomposition* of Σ an homeomorphism

$$\Phi : \Sigma \rightarrow \left(\bigcup_{i \in I} T_i \cup \bigcup_{j \in J} A_j \right) / \varphi$$

where the T_i 's are copies of T , the A_j 's are copies of A and φ is a collection of homeomorphisms reversing the orientation between boundary components of the T_i 's and the A_j 's. We ask that these homeomorphisms reduce to




Figure 2: Examples of elementary multicurves

either the identity or the complex conjugation via the identifications of the boundary components with S^1 , so that they preserve the subsets W_n . We also ask that each boundary component of a copy of T is glued to some copy of A . In that way, the components of the boundary of Σ correspond to some copies of A .

We associate to a pants decomposition of Σ the following graph G . Its vertex set is $I \cup \pi_0(\partial\Sigma)$. Edges are indexed by J in such a way that the edge associated to the annulus A_j connects the copies of T or the boundary component of Σ to which it is attached. It happens that the vertices in I are trivalent and the vertices in $\pi_0(\partial\Sigma)$ are univalent. An edge is said *internal* if it connects trivalent vertices, otherwise it is said *external*.

Consider now a surface with a pants decomposition with graph G . We will call *Dehn parameter* a pair (m, t) where m is a map from the edges of G to $\mathbb{N}$ and t is a map from the edges of G to $\mathbb{Z}$ satisfying the following conditions:

- if j_1, j_2, j_3 are three edges incoming to the same vertex, then $m_{j_1} + m_{j_2} + m_{j_3}$ is even.
- for all edges j , if $m_j = 0$ then $t_j \geq 0$.

- all external edges j satisfy $m_j = 0$.

Given such a Dehn parameter, one can construct a multicurve on Σ by gluing elementary multicurves in the following way: define $C(m, t)$ as the union $\bigcup_i C(m_1^i, m_2^i, m_3^i) \cup \bigcup_j A(m_j, t_j)/\varphi$. In this expression, m_1^i, m_2^i, m_3^i are the values of m at edges adjacent to i .

The classification theorem of Dehn is the following

Theorem 2.1. *Let Σ be a surface with pants decomposition. The map sending a Dehn parameter (m, t) to the multicurve $C(m, t)$ is a bijection between the set of Dehn parameters and the set of isotopy classes of multicurves in Σ .*

We refer to [FLP] for the proof. In the sequel we only need to know that each multicurve of Σ is isotopic to some $C(m, t)$. The fact that $C(m, t)$ is isotopic to $C(m', t')$ only if $m = m'$ and $t = t'$ is a consequence of our results.

3 Moduli spaces and torus actions

Let Σ be an oriented compact surface with boundary. Let $\mathcal{M}(\Sigma)$ be the set of isomorphism classes of pairs (E, θ) where E is a flat Hermitian bundle of rank 2 over Σ and θ is a flat unitary section of $\wedge^2 E$. An isomorphism between two pairs is an isomorphism of flat Hermitian bundles commuting with the volume sections. The holonomy representation of the fundamental group π induces an isomorphism

$$\mathcal{M}(\Sigma) \rightarrow \text{Hom}(\pi, \text{SU}(2))/\text{SU}(2)$$

To any pants decomposition is associated a torus action on a dense open subset of this moduli space together with a set of invariant functions which separate the orbits.

Let C be a simple curve (embedded circle) of Σ . Choose an orientation of C and a base point $x \in C$. Then for any $(E, \theta) \in \mathcal{M}(\Sigma)$, the holonomy at x of $E|_C$ is a unitary automorphism g_x of E_x preserving θ_x . So the trace of g_x belongs to $[-2, 2]$. We set

$$a_C(E, \theta) := \arccos\left(\frac{1}{2}\text{tr}(g_x)\right) \in [0, \pi].$$

This define a function a_C on the moduli space $\mathcal{M}(\Sigma)$ which does not depend on the choices of the base point and the orientation.

Let us introduce a circle action on the subset of $\mathcal{M}(\Sigma)$ consisting of bundles with a non-central holonomy along C . Let $t \in \mathbb{T} := \mathbb{R}/\mathbb{Z}$. To define $t.(E, \theta)$, we assume that C is oriented. Then $E|_C$ is the direct sum of two subbundles $E^\pm$, whose fibers are the eigenspaces of the holonomy along C with eigenvalues $\exp(\pm ia_C(E))$. Let R_t be the automorphism of $E|_C$ which acts on $E^\pm$ by multiplication by $\exp(\pm 2i\pi t)$.

Next consider the surface Σ' obtained from Σ by cutting out C . Denote by π the projection from Σ' to Σ . Let C_+ and C_- be the boundary components of Σ' such that π restricts to diffeomorphisms $\pi_\pm : C_\pm \rightarrow C$ and which respectively preserves and reverses the orientation. Then $t.(E, \theta)$ is the quotient of π^*E under the identification

$$u \sim \pi_-^* R_t (\pi_+^*)^{-1}(u), \quad u \in \pi^* E|_{C_+} = \pi_+^*(E|_C)$$

This definition does not depend on the orientation of C .

Let $\mathcal{C}$ be a set of simple disjoint curves. Then the corresponding actions commute and we obtain an action of the torus $\mathbb{T}^{\mathcal{C}}$ on the set $\mathcal{M}^\circ(\Sigma)$ of bundles with non-central holonomy along any curve of $\mathcal{C}$. Assume that the curves of $\mathcal{C}$ cut Σ into trinions. As in section 2 we define a graph G whose edges are naturally indexed by $J := \mathcal{C} \cup \pi_0(\partial\Sigma)$. Introduce the map

$$a : \mathcal{M}(\Sigma) \rightarrow \mathbb{R}^J$$

whose coordinates are the a_C 's.

Theorem 3.1. *The image of a is the polyhedron Δ consisting of the $(\alpha_j) \in \mathbb{R}^J$ such that for any trivalent vertex v of G*

$$|\alpha_i - \alpha_j| \leq \alpha_k \leq \min(\alpha_i + \alpha_j, 2\pi - (\alpha_i + \alpha_j))$$

if i, j and k are the edges incident to v . The fibers of the restriction of a to $\mathcal{M}^\circ(\Sigma)$ are the orbits of the action of $\mathbb{T}^{\mathcal{C}}$. Furthermore the action is locally free on the preimage of the interior of Δ .

This theorem has been proved in [JW94] under the assumption that Σ has no boundary. There is no difficulty to generalize to the case where $\partial\Sigma \neq \emptyset$. One shows first the result for a pair of pants Σ , cf. proposition 3.1 of [JW94]. The general case follows by analyzing how one can paste flat $SU(2)$ bundles on the trinions cut out by $\mathcal{C}$ to a global bundle on Σ .

When Σ has no boundary, the open set of irreducible classes (E, θ) is naturally a symplectic manifold, as we already mentioned it in section 1.3. Then for any simple curve C , the function $a_C(E)$ is a moment of the circle action corresponding to C , see [JW94] proposition 5.4.

It is easily seen that Δ has a non-empty interior. Furthermore the restrictions to $\text{Int}(\Delta)$ of the functions

$$\exp\left(i \sum_{j \in J} \ell_j a_j\right), \quad (\ell_j) \in \mathbb{Z}^J$$

are linearly independent. This will be used in the proof of theorem 5.3. The fact that Δ has a non-empty interior will also ensure that some Fourier coefficients do not vanish, cf. lemma 4.3.

4 Fourier decomposition

We are concern in this part on the Fourier decomposition on some trace functions defined on the previous moduli spaces. Consider a disjoint union Γ of unoriented circles and a map $h : \Gamma \rightarrow \Sigma$. Then define the following function on $\mathcal{M}(\Sigma)$

$$T_h(E, \theta) = \prod_{\gamma \in \pi_0(\Gamma)} (-\text{tr}(g_\gamma))$$

where g_γ is the holonomy of E along γ . Let $\mathcal{C}$ be a finite family of disjoint simple curves. Then for any $k \in \mathbb{Z}^{\mathcal{C}}$, the k -th isotype of T_h for the action of the torus $\mathbb{T}^{\mathcal{C}}$ is the function

$$\Pi_k(T_h)(x) = \int_{\mathbb{T}^{\mathcal{C}}} T_h(t.x) e^{-2\pi i \langle t, k \rangle} dt$$

The first step in the computation of $\Pi_k(T_h)$ is to extend the definition of $T_h(E, \theta)$ to the case where Γ is a disjoint union of circles and closed intervals.

4.1 Generalized holonomy trace

Let us start with an algebraic preliminary. Consider a finite family $(V_i)_{i \in I}$ of vector spaces. Let us define the tensor product $\otimes_{i \in I} V_i$ assuming that the V_i are odd superspaces. First, for any bijections σ and σ' from $\{1, \dots, n\}$ to I , one has a commutation map

$$c_{\sigma, \sigma'} : V_{\sigma(1)} \otimes \dots \otimes V_{\sigma(n)} \rightarrow V_{\sigma'(1)} \otimes \dots \otimes V_{\sigma'(n)}$$

sending $v_1 \otimes \dots \otimes v_n$ to $(-1)^{\epsilon(\alpha)} v_{\alpha(1)} \otimes \dots \otimes v_{\alpha(n)}$ where $\alpha = \sigma' \circ \sigma^{-1}$. Since $c_{\sigma', \sigma''} \circ c_{\sigma, \sigma'} = c_{\sigma, \sigma''}$, one can identify coherently these tensor products by

taking the projective limit of this system, which defines the unordered tensor product. More precisely, we define $\otimes_{i \in I} V_i$ as the subspace of the product

$$\prod_{\sigma \in \text{bij}(\{1, \dots, n\}, I)} (V_{\sigma(1)} \otimes \dots \otimes V_{\sigma(n)})$$

consisting of the families (v_σ) such that $c_{\sigma, \sigma'}(v_\sigma) = v_{\sigma'}$.

Let $E \rightarrow \Sigma$ be a flat vector bundle of rank 2 and θ a non-vanishing flat section of $\wedge^2 E$. Observe that each fiber E_x has a complex symplectic product ω_x defined in such a way that for all $u, v \in E_x$,

$$u \wedge v = \theta_x \Rightarrow \omega_x(u, v) = 1.$$

Consider a 1-dimensional compact manifold Γ and a continuous function $h : \Gamma \rightarrow \Sigma$. We will define the generalized holonomy trace of h as a vector

$$T_h(E, \theta) \in \bigotimes_{p \in \partial \Gamma} E_{h(p)}$$

If Γ is an interval, we choose an orientation of it and denote by p and q the source and target point of Γ . Then the holonomy along Γ is an homomorphism

$$A \in \text{Hom}(E_{h(p)}, E_{h(q)}) \simeq E_{h(q)} \otimes E_{h(p)}^*.$$

Identifying $E_{h(p)}$ with $E_{h(p)}^*$ by the map sending u to $\omega_{h(p)}(u, \cdot)$, we obtain a vector $A_b \in E_{h(q)} \otimes E_{h(p)}$. We set

$$T_h(E, \theta) := A_b.$$

The important point is that this definition does not depend of the orientation if we use the previous identification between $E_{h(p)} \otimes E_{h(q)}$ and $E_{h(q)} \otimes E_{h(p)}$. Indeed with this identification, we have that $(A^{-1})_b = A_b$. This is easily deduced from the fact that A is a linear symplectomorphism. If Γ is a circle, we define $T_h(E, \theta)$ as previously as the opposite of the trace of the holonomy of h (the super-trace). Finally, if Γ has several components (Γ_i) , we define $T_h(E, \theta)$ as the tensor product of the $T_{h|_{\Gamma_i}}(E, \theta)$.

Now let $h' : \Gamma' \rightarrow \Sigma$ be a continuous map and p and q be two distinct points in $\partial \Gamma'$ such that $h'(p) = h'(q)$. Then by identifying p with q , we get a compact 1-manifold Γ with a map $h : \Gamma \rightarrow \Sigma$. It is easily checked that

$$T_h(E, \theta) = C(T_{h'}(E, \theta))$$

where C is the contraction

$$\bigotimes_{r \in \partial \Gamma'} E_{h(r)} \longrightarrow \bigotimes_{r \in \partial \Gamma' \setminus \{p, q\}} E_{h(r)}$$

sending $u \otimes v \otimes w$ to $u \cdot \omega(v, w)$ if $w \in E_{h(p)}$, $v \in E_{h(q)}$ and $u \in \bigotimes_{r \in \partial \Gamma' \setminus \{p, q\}} E_{h(r)}$.

4.2 The basic computation

In this part we consider the case of a single simple curve C of Σ with its associated circle action. Let Γ be a finite disjoint union of circles and $h : \Gamma \rightarrow \Sigma$ be a map intersecting C transversally.

Let Γ' be the compact 1-manifold obtained by cutting out the points p of Γ such that $h(p)$ is on C and let h' be the obvious map $\Gamma' \rightarrow \Sigma$. Then it follows from the considerations of section 4.1 that $T_h(E, \theta)$ is the contraction of the generalized trace holonomy $T_{h'}(E, \theta)$ by the linear map

$$C : \bigotimes_{p \in \partial \Gamma'} E_{h'(p)} \rightarrow \mathbb{C}$$

defined as follows. For any $p \in h^{-1}(C)$, let $j_1(p)$ and $j_2(p)$ be the corresponding boundary points of Γ' , so that

$$\bigotimes_{p \in \partial \Gamma'} E_{h'(p)} = \bigotimes_{p \in h^{-1}(C)} E_{h(p)}^{\otimes \{j_1(p), j_2(p)\}}$$

Then C is the map sending $\bigotimes_{p \in h^{-1}(C)} (v_{p,1} \otimes v_{p,2})$ to $\prod_{p \in h^{-1}(C)} \omega_{h(p)}(v_{p,1}, v_{p,2})$.

We will compute $\Pi_k(T_h)$ in terms of these generalized holonomy traces. Assume that C is oriented and pick (E, θ) such that its holonomy along C is not central. For any $p \in h^{-1}(C)$, we choose two unitary eigenvectors $e_{\pm}^p \in E_{h(p)}$ of the holonomy along C with corresponding eigenvalues $\exp(\pm i a_C(E))$ and such that

$$\theta_{h(p)} = e_+^p \wedge e_-^p.$$

Assume also that $j_1(p)$ and $j_2(p)$ are chosen in such a way that the tangent vector of Γ oriented from j_1 to j_2 followed with the oriented tangent vector of C form a direct basis in Σ .

Lemma 4.1. *Let $k \in \mathbb{N}$. If k is bigger than the cardinal of $h^{-1}(C)$, then $\Pi_k(T_h)$ and $\Pi_{-k}(T_h)$ vanish. If k is equal to the cardinal of $h^{-1}(C)$, then*

$$\Pi_{\pm k}(T_h)(E, \theta) = C_{\pm}(T_{h'}(E, \theta)),$$

with $C_\pm$ the linear map from $\bigotimes_{p \in h^{-1}(C)} E_{h(p)}^{\otimes \{j_1(p), j_2(p)\}}$ to $\mathbb{C}$ given by

$$C_\pm \left(\bigotimes_{p \in h^{-1}(C)} (v_{p,1} \otimes v_{p,2}) \right) = \prod_{p \in h^{-1}(C)} \langle v_{p,1}, e_\pm^p \rangle \langle v_{p,2}, e_\mp^p \rangle$$

where the bracket denotes the scalar product. If C does not intersect h , $C_\pm$ is the identity map.

Proof. Assume first that $h^{-1}(C)$ is reduced to a point. Then the proof is based on the simple observation that

$$T_h(t.(E, \theta)) = C_t(T_{h'}(E, \theta)), \quad \forall t \in \mathbb{T} \quad (4)$$

where C_t is the linear map from $E_{h(p)}^{\otimes \{j_1(p), j_2(p)\}}$ to $\mathbb{C}$ given by

$$C_t(v_1 \otimes v_2) = \omega_{h(p)}(R_t v_1, v_2)$$

with R_t the automorphism of E_C entering in the definition of the circle action. Next a straightforward computation leads to

$$\omega_{h(p)}(R_t v_1, v_2) = e^{2i\pi t} \langle v_1, e_+^p \rangle \langle v_2, e_-^p \rangle + e^{-2i\pi t} \langle v_1, e_-^p \rangle \langle v_2, e_+^p \rangle$$

which proves the result, taking into account that the Hermitian product is $\mathbb{C}$ -linear on the left. In the case where $h^{-1}(C)$ consists of several points, (4) is still satisfied with the contraction

$$C_t \left(\bigotimes_{p \in h^{-1}(C)} (v_{p,1} \otimes v_{p,2}) \right) = \prod_{p \in h^{-1}(C)} \omega_{h(p)}(R_t v_{p,1}, v_{p,2}).$$

Replacing each term in the product by the previous formula leads to the result. $\square$

4.3 Fractional Dehn twist

Consider again a curve C on Σ and a map $h : \Gamma \rightarrow \Sigma$ which is transverse to C in k points. One defines the *fractional Dehn twist* of h along C of order ℓ/k in the following way: let Γ' be the 1-manifold Γ obtained by cutting out the points p such that $h(p)$ is on C . Let us orient C , then for any p in $P = C \cap h(\Gamma)$, one defines two points $j_1(p)$ and $j_2(p)$ in $\partial\Gamma'$ with the same convention as in the section 4.2. Moreover, the orientation of C provides P with a cyclic order, so $\mathbb{Z}/k\mathbb{Z}$ acts on P . One sets

$$\Gamma_{\ell/k} = \Gamma' \cup \coprod_{p \in P} [0, 1]_p / \sim$$

where we identify 0_p with $j_1(p)$ and 1_p with $j_2(p + \ell)$.

Choose a parameterization $\alpha : S^1 \rightarrow C$ respecting the orientation such that P corresponds to the set of k -th roots of unity. We define the map $h_{\ell/k} : \Gamma_{\ell/k} \rightarrow \Sigma$ as being equal to h on Γ' and such that for all $t \in [0, 1]$ and $p \in P$ one has

$$h_{\ell/k}(t_p) = \alpha(\exp(\frac{2i\pi\ell}{k}t_p) \cdot \alpha^{-1}(p)).$$

The manifold $\Gamma_{\ell/k}$ and the map $h_{\ell/k}$ do not depend up to homotopy on the orientation of C nor on the parameterization α .

Lemma 4.2. *One has $\Pi_{\pm k}(T_{h_{\ell/k}}) = (-1)^{\ell(k-1)} e^{\pm i\ell a_C} \Pi_{\pm k}(T_h)$*

Proof. This is an easy application of the lemma 4.1 using the explicit description of the fractional Dehn twist above. More precisely, denote by $H_{\ell/k}$ the restriction of $h_{\ell/k}$ to $\coprod_{i=1}^k [0, 1]_i$ and choose a parameterization $\alpha : S^1 \rightarrow C$ as above. Given a bundle E in $\mathcal{M}(\Sigma)$ one can suppose that it is trivialized on P in such a way that the holonomy from p to $p + 1$ is

$$\begin{bmatrix} \exp(2i\pi a_C/k) & 0 \\ 0 & \exp(-2i\pi a_C/k) \end{bmatrix}.$$

Then $H_{\ell/k}$ and $H_{0/k}$ differ by the factor $\exp(2i\pi a_C/k)^{k\ell}$ and the sign of the permutation sending x to $x + \ell$ in $\mathbb{Z}/k\mathbb{Z}$. $\square$

4.4 Non vanishing of some isotypes

Consider a pants decomposition of Σ in the sense of section 2 with graph G . Let $S^1 \times \{1/2\}$ be the core of the standard annulus $S^1 \times [0, 1]$. For each internal edges of G , we consider the core of the corresponding annulus and the circle action it generates. This defines a torus action of $\mathbb{T}^{\mathcal{C}}$ where $\mathcal{C}$ is the set of internal edges.

Consider the multicurve $C(m, t)$ with Dehn coordinates (m, t) . By the first part of Lemma 4.1, the k -th isotype of $T_{C(m, t)}$ vanishes if $|k_j| > m_j$ for some internal edge j .

Lemma 4.3. *Let $k \in \mathbb{Z}^{\mathcal{C}}$ be such that $|k_j| = m_j$ for all $j \in \mathcal{C}$. Then the function $\Pi_k(T_{C(m, t)})$ does not vanish on $a^{-1}(\text{Int}(\Delta))$, where a is the map introduced in section 3 and Δ is the image of a .*

Proof. By lemma 4.2, one may assume that t vanishes. One can compute $\Pi_k(T_{C(m, 0)})$ in exactly the same way as we did in section 4.2. The cut manifold Γ' is now a disjoint union of closed intervals with a map $h' : \Gamma' \rightarrow \Sigma$.

For any oriented curve C in $\mathcal{C}$ and point p of C , one considers a unitary eigenvector e of E_p for the holonomy along C whose eigenvalue has positive imaginary part. To any point p in $\partial\Gamma'$ such that $h'(p) \in C$, one associates the vector e_+^p (resp. e_-^p) corresponding to the orientation of C going to the left from p (resp. to the right).

By generalizing lemma 4.1, one obtains that

$$\Pi_k(T_h)(E, \theta) = C_k(T_{h'}(E, \theta)),$$

with C_k a linear map from $\bigotimes_{p \in \partial\Gamma'} E_{h'(p)}$ to $\mathbb{C}$ of the form

$$C_k(\bigotimes_{p \in \partial\Gamma'} v_p) = \prod_{p \in \partial\Gamma'} \langle v_p, e_{\epsilon(k,p)}^p \rangle \mod \mathbb{T}$$

where $\epsilon(k, p)$ is the sign of k_j if p belongs to $j \in \mathcal{C}$. Since $T_{h'}(E, \theta)$ is the tensor product of the $T_{h'|_\gamma}(E, \theta)$ where γ runs over the components of Γ' , one can rewrite the previous product as

$$\Pi_k(T_h)(E, \theta) = \prod_{\gamma \in \pi_0(\Gamma')} C_k^\gamma(T_{h'|_\gamma}(E, \theta)) \mod \mathbb{T}$$

with $C_k^\gamma(v_p \otimes v_q) = \langle v_p, e_{\epsilon(k,p)}^p \rangle \langle v_q, e_{\epsilon(k,q)}^q \rangle$ if the source and target of γ are p and q .

To prove the lemma, it suffices to show that the factors $C_k^\gamma(T_{h'|_\gamma}(E, \theta))$ do not vanish when $a(E, \theta)$ belongs to the interior of Δ . There are two cases to consider, according to whether the endpoints of γ belong to the same separating curve or to different ones, cf. for example $C(2, 0, 0)$ and $C(1, 1, 0)$.

In the second case, consider the pair of pants containing γ and let j_1, j_2 and j_3 be the bounding curves with the induced orientation. Assume that the endpoints p and q of γ belongs to j_1 and j_2 respectively. If $C_k^\gamma(T_{h'|_\gamma}(E, \theta))$ vanishes, one deduces from the definition of $T_{h'|_\gamma}(E, \theta)$ that the holonomy along the loop $h'|_\gamma$ sends the decomposition $\mathbb{C}e_+^p \oplus \mathbb{C}e_-^p$ to the decomposition $\mathbb{C}e_+^q \oplus \mathbb{C}e_-^q$, permuting possibly the summands. Choosing p as a base point, one obtains that the holonomies along j_1 and the concatenation $\gamma^{-1}j_2\gamma$ commute. Observe furthermore that j_3^{-1} is freely isotopic to $\gamma^{-1}j_2\gamma j_1$. It follows that for some $\varepsilon_i \in \{1, -1\}$, the three eigenvalues $\exp(\varepsilon_1 i a_{j_1}(E)), \exp(\varepsilon_2 i a_{j_2}(E)), \exp(\varepsilon_3 i a_{j_3}(E))$ have product equal to 1. This implies that

$$\varepsilon_1 a_{j_1}(E) + \varepsilon_2 a_{j_2}(E) + \varepsilon_3 a_{j_3}(E) \equiv 0 \mod 2\pi\mathbb{Z}$$

This formula is only satisfied if $a(E, \theta)$ belongs to the boundary of Δ .

In the first case, $h'|_\gamma$ connects two points p and q of the same boundary circle j_1 by going around another boundary circle j_2 . Since the neighborhoods of the endpoints of γ are on the same side of j_1 , the eigenvalues of e_p and e_q are the same. Using this, one shows that if $C_k^\gamma(T_{h'|_\gamma}(E, \theta))$ vanishes, then the holonomy along a path joining j_1 to j_2 preserves the decomposition into eigenspaces. In other words, we are again in the second case. $\square$

5 Consequences

5.1 Geometric intersection numbers

The *geometric intersection number* of two isotopy classes of multicurves ξ and η is the minimal number of intersection points of a representant of ξ with a representant of η .

If ξ has only one connected component, it generates a circle action on the $\mathcal{M}(\Sigma)$. One may characterize the geometric intersection number of ξ and η in terms of this action and the trace function of η .

Theorem 5.1. *For any isotopy classes ξ and η of a curve and a multicurve respectively, the geometric intersection number of ξ and η is the biggest k such that the k -th isotype of the trace function of η with respect to the circle action generated by ξ does not vanish.*

Proof. If k is larger than the geometric intersection number, then the k -th isotype vanishes by Lemma 4.1. Conversely, assume that ξ does not bound a disc and is not parallel to a boundary component, otherwise the result is trivially satisfied. Then there is a pants decomposition with a separating curve C representing ξ . By Theorem 2.1, η may be represented by a Dehn multicurve $C(m, t)$. Then lemma 4.3 shows that the k -isotype of $T_{C(m, t)}$ does not vanish if k is the number of intersection points of $C(m, t)$ with C . $\square$

The proof shows also that the intersection number is realized when the representant of ξ is a curve of a pants decomposition and the representant of η is one of the associated Dehn curves, a well-known result.

The following corollary has been proved in [Gol86]. Denote by $\mathcal{M}^s(\Sigma)$ the subset of $\mathcal{M}(\Sigma)$ consisting of the irreducible bundles.

Corollary 5.2. *Assume that Σ is closed so that $\mathcal{M}^s(\Sigma)$ is a symplectic manifold. Let ξ be a curve and η a multicurve in Σ . Then the Poisson*

bracket of the trace functions of ξ and η vanishes if and only if ξ and η admit non-intersecting representants.

This follows from the fact that the function $\arccos(-\frac{1}{2}T_\xi)$ is a moment of the circle action generated by ξ .

5.2 Independence of trace functions

One can now prove the following theorem:

Theorem 5.3. *The functions T_ξ , where ξ runs over isotopy classes of multicurves in Σ , are linearly independent.*

Since the isomorphism between $\mathcal{M}(\Sigma)$ and $\text{Hom}(\pi, \text{SU}(2))/\text{SU}(2)$ identifies the function T_ξ with the trace function $f_{[\xi], \text{SU}(2)}$, we deduce theorem 1.1 for the group $\text{SU}(2)$. Consider the map j

$$\text{Hom}(\pi, \text{SU}(2))/\text{SU}(2) \rightarrow \text{Hom}(\pi, \text{Sl}(2, \mathbb{C}))/\text{Sl}(2, \mathbb{C})$$

Then $j^* f_{\gamma, \text{Sl}(2, \mathbb{C})} = f_{\gamma, \text{SU}(2)}$. So the independence of the $f_{\gamma, \text{SU}(2)}$ implies the independence of the $f_{\gamma, \text{Sl}(2, \mathbb{C})}$, which proves theorem 1.1 for the group $\text{Sl}(2, \mathbb{C})$.

Proof. Consider a pants decomposition of Σ with associated graph G and the action of the torus $\mathbb{T}^\mathcal{C}$ as in section 4.4. Let $J = \mathcal{C} \cup \pi_0(\Sigma)$ be the set of edges of G . By theorem 2.1, one may suppose that ξ runs over the multicurves $C(m, t)$ for admissible maps $m : J \rightarrow \mathbb{N}$ and $t : J \rightarrow \mathbb{Z}$.

Let β be a vanishing linear combination of the T_ξ 's. Let $\beta = \sum_m \beta_m$ be its decomposition with respect to the multi-degree m . Write

$$\beta_m = \sum_t \lambda_{m,t} T_{C(m,t)}.$$

Consider an arbitrary order on J and the corresponding lexicographical order on maps m . Let M be the maximal m with a non-vanishing family of coefficients $(\lambda_{m,t})_t$. By lemma 4.1, for $m < M$ one has $\Pi_M(\beta_m) = 0$. Hence $\Pi_M(\beta_M) = \Pi_M(\beta) = 0$. Furthermore, by lemma 4.2,

$$\begin{aligned} \Pi_M(\beta_M) &= \sum_t \lambda_{M,t} \Pi_M(T_{C(M,t)}) \\ &= \left(\sum_t \lambda_{M,t} \prod_{j \in J} \varphi_{M_j, t_j}(a_j) \right) \Pi_M(T_{C(M,0)}) \end{aligned}$$

where for any integers k, ℓ and real α ,

$$\varphi_{k,\ell}(\alpha) = \begin{cases} (-1)^{\ell(k-1)} \exp(i\ell\alpha) & \text{if } k \neq 0 \\ (2 \cos(\alpha))^\ell & \text{otherwise} \end{cases}$$

By lemma 4.3, the function $\Pi_M(T_{C(M,0)})$ does not vanish on $a^{-1}(\text{Int } \Delta)$. Furthermore, by the remark after theorem 3.1, the functions

$$\prod_{j \in J} \varphi_{M_j, t_j}(a_j)$$

where t runs over the maps $J \rightarrow \mathbb{Z}$ such that (M, t) are Dehn coordinates, are linearly independent over $a^{-1}(\text{Int } \Delta)$. So the coefficients $\lambda_{M,t}$ vanish, contradicting the maximality of M . $\square$

6 On the character variety

Denote by V the vector space $\mathbb{C}^2$ and by G the group $\text{Sl}(2, \mathbb{C}) \subset \text{End}(V)$. Let π be a finitely generated group. Choosing generators $t_1, \dots, t_n$ of π , we identify the set $\text{Hom}(\pi, G)$ of morphisms with a closed algebraic subset of $(\text{End } V)^n$ by sending the morphism ρ to $(\rho(t_1), \dots, \rho(t_n))$. This endows $\text{Hom}(\pi, G)$ with a structure of affine variety. Its coordinate ring $\mathbb{C}[\text{Hom}(\pi, G)]$ is the quotient of $\mathbb{C}[(\text{End } V)^n]$ by the ideal of polynomial functions vanishing on $\text{Hom}(\pi, G)$.

The action by conjugation of G on $\text{Hom}(\pi, G)$ is regular. By Hilbert theorem, the ring of invariant functions is finitely generated. By definition $\mathbb{C}[\text{Hom}(\pi, G)]^G$ is the coordinate ring of the character variety.

One may naturally identify this coordinate ring with a subspace of the space of complex valued functions on the quotient $\text{Hom}(\pi, G)/G$. By the following theorem, this subspace is the space $\mathcal{T}(\pi, G)$ of trace functions.

Theorem 6.1. *The ring $\mathbb{C}[\text{Hom}(\pi, G)]^G$ is generated as a vector space by the functions χ_t*

$$\chi_t([\rho]) = -\text{tr}(\rho(t)), \quad \rho \in \text{Hom}(\pi, G)$$

where t runs over π .

We provide an elementary proof of this well-known result.

Proof. Because of the trace relation (1), it suffices to prove that the ring of regular invariant functions is generated by the χ_t 's as an algebra. Then by

averaging the action of the compact subgroup $\mathrm{SU}(2) \subset G$, we get that any invariant regular function is represented by an invariant polynomial function $P \in \mathbb{C}[(\mathrm{End} V)^n]^G$.

The symplectic form of V induce an isomorphism between V and V^* and consequently an isomorphism between $\mathrm{End} V$ and $\mathrm{End} V^*$. By composing with the transposition map, we define an equivariant isomorphism $a \rightarrow a^*$ of $\mathrm{End}(V)$. We will prove that $\mathbb{C}[(\mathrm{End} V)^n]^G$ is generated as an algebra by the functions

$$(a_1, \dots, a_n) \rightarrow \mathrm{tr}(w(a)) \quad (5)$$

where $w(a)$ runs over the words in the letters $a_1, a_1^*, \dots, a_n, a_n^*$. Since the elements of G satisfy $a^{-1} = a^*$, this will end the proof.

Let $W = \mathrm{End}(V)$. For any n -tuple $(d_1, \dots, d_n)$ of non negative integers, the space of polynomials on $W^{\oplus n}$ which are homogeneous of degree d_i on the i -th variable is isomorphic with $\otimes_i \mathrm{Sym}^{d_i}(W_i^*)$. So

$$\mathbb{C}[W^{\oplus n}] = \bigoplus_{d \in \mathbb{N}^n} \bigotimes_{i=1, \dots, n} \mathrm{Sym}^{d_i}(W^*) \subset \bigoplus_{d \in \mathbb{N}^n} \bigotimes_{i=1, \dots, n} (W^*)^{\otimes d_i} =: E$$

A right inverse of this inclusion is the following map from E to $\mathbb{C}[W^{\oplus n}]$

$$\bigotimes_{i=1, \dots, n} (\ell_i^1 \otimes \dots \otimes \ell_i^{d_i}) \rightarrow P(a_1, \dots, a_n) = \prod_{\substack{i=1, \dots, n; \\ j=1, \dots, d_i}} \ell_i^j(a_i)$$

Observe that the invariant subspace of E is sent onto $\mathbb{C}[W^{\oplus n}]^G$. Using again the identification of V with V^* , we have that $W \simeq V \otimes V^* \simeq V^* \otimes V^*$. By lemma 6.2, the invariant subspace of $(V^*)^{\otimes 2m} \simeq (V^{\otimes 2m})^*$ is generated by the maps

$$x_1 \otimes \dots \otimes x_{2m} \rightarrow \omega(x_{\sigma(1)}, x_{\sigma(2)}) \dots \omega(x_{\sigma(2m-1)}, x_{\sigma(2m)})$$

where σ runs over the permutation of $\{1, \dots, 2m\}$. We deduce the invariants subspace of E and then that the functions (5) generate $\mathbb{C}[W^{\oplus n}]^G$. $\square$

Denote by Mult_n the space of multilinear maps from $V^{\times n}$ to $\mathbb{C}$ and by Sym_n the subspace of symmetric maps. For any non-negative integer $k \leq n$ with the same parity as n , let $\mathrm{Mult}_{n,k}$ be the subspace of Mult_n generated by the maps

$$P(x_{\sigma(1)}, \dots, x_{\sigma(k)}) \omega(x_{\sigma(k+1)}, x_{\sigma(k+2)}) \dots \omega(x_{\sigma(n-1)}, x_{\sigma(n)})$$

where σ ranges over the permutations of $\{1, \dots, n\}$ and P over Sym_k . Recall that the spaces Sym_n are the irreducible representations of G .

Lemma 6.2. *The decomposition into isotypical subspaces of Mult_n is*

$$\text{Mult}_n = \bigoplus_{\substack{0 \leq k \leq n, \\ k \equiv n \pmod{2}}} \text{Mult}_{n,k}$$

Proof. $\text{Mult}_{n,k}$ is clearly a subspace of the k -th isotypical component of Mult_n . Since Mult_n is the direct sum of its isotypical components, we just have to show that the spaces $\text{Mult}_{n,k}$ generates Mult_n . This may be proved by induction over n by using that any multilinear map symmetric with respect to the $(n-1)$ first arguments is of the form

$$L(x_1, \dots, x_n) = M(x_1, \dots, x_n) + \sum_{i=1}^n \omega(x_i, x_n) N(x_1, \dots, \widehat{x_i}, \dots, x_{n-1})$$

with $M \in \text{Sym}_n$ and $N \in \text{Sym}_{n-2}$. To show this last fact, observe that the map sending (M, N) to L is an isomorphism from $\text{Sym}_n \oplus \text{Sym}_{n-2}$ onto the subspace of Mult_n consisting of maps symmetric in the $(n-1)$ first arguments. Indeed, this morphism is injective by G -equivariance and we conclude by counting dimensions. $\square$

References

- [And06] Jorgen E. Andersen. Asymptotic faithfulness of the quantum $\text{SU}(n)$ representations of the mapping class groups. *Ann. of Math.*, 163(2):347–368, 2006.
- [AB83] Michael F. Atiyah and Raoul Bott. The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A*, 308(1505):523–615, 1983.
- [BHMV95] C. Blanchet, N. Habegger, G. Masbaum and P. Vogel. Topological quantum field theories derived from the Kauffman bracket. *Topology*, 34:883–927, 1995.
- [Bul97] Doug Bullock. Rings of $\text{SL}_2(\mathbf{C})$ -characters and the Kauffman bracket skein module. *Comment. Math. Helv.*, 72(4):521–542, 1997.
- [BFK99] D. Bullock, C. Frohman and J. Kania-Bartoszyńska. Understanding the Kauffman bracket skein module. *J. Knot Theory Ramifications*, 8(3):265–277, 1999.

- [FLP] A. Fathi, F. Laudenbach and V. Poenaru. Travaux de Thurston sur les surfaces. *Asterisque*, 1979.
- [FWW02] M. H. Freedman, K. Walker and Z. Wang. Quantum $SU(2)$ asymptotically detects mapping class group modulo center. *Geom. Topol.*, 6:523–539, 2002.
- [Gol84] William M. Goldman. The symplectic nature of fundamental groups of surfaces. *Adv. in Math.*, 54(2):200–225, 1984.
- [Gol86] William M. Goldman. Invariant functions on Lie groups and hamiltonian flows of surface group representations. *Invent. Math.*, 85(2):263–302, 1986.
- [GM93] F. Gonzalez-Acuña and J. Montesinos. On the character variety of group representations in $SL_2(\mathbb{C})$ and $SL_2(\mathbb{C})$. *Math. Z.*, 214:627–652, 1993.
- [JW94] Lisa C. Jeffrey and Jonathan Weitsman. Bohr-Sommerfeld orbits in the moduli space of flat connections and the Verlinde dimension formula. *Comm. Math. Physics*, 150(3):593–630, 1992.
- [JW94] Lisa C. Jeffrey and Jonathan Weitsman. Toric structures on the moduli space of flat connections on a Riemann surface: volumes and the moment map. *Adv. Math.*, 106(2):151–168, 1994.
- [MN08] Julien Marche and Majid Narimannejad. Some asymptotics of topological quantum field theory via skein theory. *Duke Math. J.*, 141(3):573–587, 2008.
- [PS00] Józef H. Przytycki and Adam S. Sikora. On skein algebras and $SL_2(\mathbb{C})$ -character varieties. *Topology*, 39(1):115–148, 2000.
- [T94] Vladimir G. Turaev. Skein quantization of Poisson algebras of loops on surfaces. *Ann. Sci. École Norm. Sup.*, 24(6):635–704, 1994.